



## Notice for Port-A Catheter

Created in 1998.06  
Reviewed in 2022.11  
English version in 2022.11

### I. Introduction :

A port-A catheter is a medical device, which is designed to facilitate access to the venous system. The size of port-A is about the size of a 10 NT dollar coin. Where port-A place you is a round hump. The device consists of a catheter which is inserted into a vein, and a reservoir which is implanted completely under the skin. It is used for long term and frequent injections via intravenous treatments. As long as the port-A is placed correctly and under adequate care, complications can be minimized. It can be set safely and efficiently.

### II. Port-A placement :

1. Doctors can determine the most appropriate method for port-A insertion. The port-A is implanted usually on the subclavian, superior vein cava, or jugular vein.
2. This is a simple operation that only require local anesthesia.
3. After surgery, X-rays will be taken according to the location of the device to confirm the correct position of the device.

### III. Post surgery wound care :

1. The Wound would usually be painful about 3-4days, you can use painkillers available as prescribed, then subside gradually. Notify the nursing staff if you feel any discomfort.
2. Wound healing takes about 5~7 days with a 2~3cm scar formation. You don't have to remove suture line. The wound covered in a dressing to protect the site and to secure the needle in position. Make sure to Keep the wound dry. Nurses will check the wound every day, performing wound care according its situation. It can be absorbed by human body.
3. Port-A can be used right away if there is no discomfort.
4. During the week after operation, don't do strenuous activity on same side of the wound to avoid port-A displacement.

### IV. Recommendations :

1. Call nurse or doctor right away if the wound gets redness, swells, burning sensation or painful.
2. Avoid doing following things on the side of the wound of body: collision, lifting heavy stuff, intense chest expansion movement, swimming, weight lifting,

- playing golf, doing taiji, combing hair, drying hair, and etc.
3. Don't rotate the limb beside the wound 360 degree or the base of port-A would rotate. If the base of port-A rotate too much, port-A will be unable to use.
  4. During the injection via port-A, do not press needles and do not let the hand cross to the opposite side of the body to avoid the needle displacement and avoid lying on the injection side when sleeping.
  5. Call nurse or doctor right away if you feel discomfort during injection.
  6. If treatment is finished, after evaluation by doctor, Port-A can be removed.
  7. The port-A catheter also needs to be cleaned up periodically to eliminate the risk of developing of blood clots.

**Artery system : once a week**

**Venous system : One to three months**

**Intra-abdominal system : once a week**

If you have any questions or suggestions, please contact us. We are very willing to help. Our contact telephone number: Taipei main branch Hospital: (02)25713760 ; Hsinchu branch: (03)5745098 ; Taitung branch: (089) 310150 ext: 311. Time of consultant: 9:00 am.~12:00 noon,2:00 pm.~5:00 pm Monday ~ Friday.

**May God Bless You**